

# The Ecological Niche Part 1: Competition

---

Michael Noonan

Biol 417: Evolutionary Ecology

1. Review
2. Niche History
3. Niche Overlap and Competition

# Review

---

Over the past week we have covered how meteorology, climate, and geography govern the physical properties of any given place on Earth. This dictates which strategies are effective/not in any environment, in turn dictating the structures of ecosystems, a phenomenon termed 'Biogeography'.

Meteorology and biogeography form the backdrop against which any evolutionary/ecological phenomenon must be considered.

Today we will continue along this line of thought and talk about the ecological 'niche'.



## Niche History

---

The field of biogeography focuses on understanding spatial patterns of ecological communities.



Biogeographers quickly realised that some species are abundant in certain locations but non-existent in others.

I.e., there are conditions where they do well, and others there they can't survive and reproduce.

In 1917, Grinnell (1917) used the word 'niche' as an umbrella term to describe the range of environments and conditions that the California Thrasher inhabits.



Source: <https://birdsoftheworld.org/>

Grinnell was interested primarily in describing the abiotic factors underpinning where a species could occur.

A species' 'Grinnellian' niche is defined by abiotic variables (e.g., temperature, precipitation, solar radiation, soil conditions, etc...).

But what about species interactions (e.g., predation, competition, symbiosis, etc...)?

The Grinnellian niche provides no framework for understanding a species' place in the environment in relation to **biotic** factors.  
(major limitation)

Ten years later, Elton (2001) extended the definition of the niche to mean a species' "... place in the biotic environment, its relations to food and enemies."

Conceptually, the Eltonian niche introduces the idea of a species' response to and effect on the environment.

Elton emphasised that species not only grow and respond to environmental conditions, but that they also change the availability and behavior of those factors as they grow. (!)


The Eltonian niche helped (evolutionary) ecologists conceptualise complicated ecological relationships that have been playing out over evolutionary timescales.

It has (and continues to) served ecologists well for nearly a century, but it has a major shortcoming:

How do we quantify a niche?


At the 1957 Cold Spring Harbor Symposia on Quantitative Biology there was a debate around the niche concept. After the conference Hutchinson (1957) wrote a paper called “Concluding Remarks”.

He started with the idea that plots of fitness versus a single environmental gradient usually look like bell-shaped curves



Source: Pianka (2000)


You can easily extend the idea to two different environmental variables



Source: Pianka (2000)




... and to three different environmental variables




Source: Pianka (2000)

... and so on for any number of axes generating an **n-dimensional hypervolume** enclosing the complete range of conditions under which it can successfully replace itself.

Volume where a species' growth rate,  $r_0$ , is  $\geq 0$  are inside the niche, volume with  $r_0 < 0$  are outside the niche.



*Calandra oryzae*, source: Birch (1953)




*Daphnia magna*, source: Hooper *et al.* (2008)

## Niche Overlap and Competition


---

Hutchinson's n-dimensional hypervolume niche identifies all the conditions under which a species can live, grow, and replace itself.

This space is called the **fundamental** niche





Most species only occupy a fraction of their fundamental niche: i.e., the **realised** niche




In trying to explain discrepancies between fundamental and realised niches Hutchinson focused on the concept of niche overlap and competition.


He argued that habitats should be fully saturated and niche overlap should not be permitted (i.e., overlap  $\rightarrow$  competitive exclusion).



## Potential scenarios:




Finding: discrepancies between fundamental and realised niches were the direct result of competitive exclusion.



Based on: Connell (1961)


Gentoo (*P. papua*) and Adélie (*P. adeliae*) penguins are ecologically similar and feed primarily on Antarctic krill (*E. superba*). Spatially distinct foraging niches permit co-existence.



Source: Pickett *et al.* (2018)



When two species of mud snails (*Hydrobia spp.*) occupy different habitats, their shell sizes are identical and they specialise on food of the same size. When they occupy the same habitat they each specialise on food of different sizes and their shell sizes differ.



Source: Fenchel & Christiansen (1977)

Problem: No explanation for niche overlap without competition.



Source: <http://www.clydesideimages.co.uk/>


Hutchinson (1957) laid out assumptions to translate the **qualitative** concept of a niche into a **quantitative** concept. Making these assumptions introduced three important limitations/properties into his framework:

1. Saturation
2. Dimensionality
3. Non-competitive interactions

Hutchinson (1957) assumed that the world is completely saturated.

Under this scenario, any niche space given is niche space lost, so competition is fierce. Are we prepared to make that assumption?


The amount of tolerable overlap will depend on resource availability.



Source: Pianka (2000)


Hutchinson (1957) defined an n-dimensional hypervolume.

Niches are usually evaluated across 1-2 axes, but overlap on one dimension doesn't mean overlap across the entire hypervolume.



Source: Pianka (2000)

In *Anolis* spp. lizards, pairs of species with high dietary overlap often exploit different structural microhabitats.



Source: Schoener (1968)

Hutchinson (1957) focused on competitive interactions and assumed any niche overlap decreased  $r_0$

... but there are many conditions where niche overlap can increase  $r_0$  (e.g., predator saturation, many eyes, etc...).



Source: <http://www.clydesideimages.co.uk/>

The concept of the niche is central to evolutionary ecology (understand a species' fundamental vs. realised niche and you can understand much of its evolutionary ecology).

Quantifying a species' entire niche space is borderline impossible, but quantifying too few dimensions can ignore important separation.

There is no perfect framework for studying niches, but in doing so we have learned a lot about how species exist and co-exist.

We will continue along this train of thought next lecture...



# References

---

- Birch, L. (1953). Experimental background to the study of the distribution and abundance of insects: li. the relation between innate capacity for increase in numbers and the abundance of three grain beetles in experimental populations. *Ecology*, 34, 712–726.
- Connell, J.H. (1961). The influence of interspecific competition and other factors on the distribution of the barnacle *chthamalus stellatus*. *Ecology*, pp. 710–723.
- Elton, C.S. (2001). *Animal ecology*. University of Chicago Press.
- Fenchel, T.M. & Christiansen, F.B. (1977). Selection and interspecific competition. In: *Measuring selection in natural populations*. Springer, pp. 477–498.
- Grinnell, J. (1917). The niche-relationships of the california thrasher. *The Auk*, 34, 427–433.
- Hooper, H.L., Connon, R., Callaghan, A., Fryer, G., Yarwood-Buchanan, S., Biggs, J., Maund, S.J., Hutchinson, T.H. & Sibly, R.M. (2008). The ecological niche of *daphnia magna* characterized using population growth rate. *Ecology*, 89, 1015–1022.
- Hutchinson, G.E. (1957). Cold spring harbor symposium on quantitative biology. *Concluding remarks*, 22, 415–427.
- Pianka, E.R. (2000). *Evolutionary Ecology*. 6th edn. Benjamin/Cummings, San Francisco.
- Pickett, E.P., Fraser, W.R., Patterson-Fraser, D.L., Cimino, M.A., Torres, L.G. & Friedlaender, A.S. (2018). Spatial niche partitioning may promote coexistence of pygoscelis penguins as climate-induced sympatry occurs. *Ecology and evolution*, 8, 9764–9778.
- Schoener, T.W. (1968). The anolis lizards of bimini: resource partitioning in a complex fauna. *Ecology*, 49, 704–726.